

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti E-Cadherin / Cadherin-1, IgG1, Clone: [5H9], Monoclonal NMB-MUB0301P

Artikelname	Mouse anti E-Cadherin / Cadherin-1, IgG1, Clone: [5H9], Monoclonal
Artikelnummer	NMB-MUB0301P
Hersteller Artikelnummer	MUB0301P
Alternativnummer	NMB-MUB0301P
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	ICC, IHC-Fr, IHC-P, WB
Spezies Reaktivität	Human
Produktbeschreibung	Cadherins constitute a family of transmembrane glycoproteins involved in Ca ²⁺ -dependent cell-cell interactions. The members of this family are differentially expressed in various tissues. They function in the maintenance of tissue integrity and morph...
Klonalität	Monoclonal
Klon-Bezeichnung	[5H9]
Isotyp	IgG1
UniProt	P12830
Puffer	Each vial contains 100 ul of 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

Quelle	5H9 is a Mouse monoclonal IgG1 antibody obtained by fusion of P3-X63-Ag 8,653 Mouse myeloma cells with spleen cells from a BALB/c Mouse immunized with affinity purified 80 kD extracellular fragments of E-cadherin derived from tryptic digestion of A-431 Hu
Formel	Each vial contains 100 ul of 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.
Anwendungsbeschreibung	5H9 is suitable for immunoblotting, immunocytochemistry and immunohistochemistry on frozen and paraffin embedded tissues. For frozen tissues we recommend fixation with 1% formaldehyde in PBS buffer containing 0.1 mM CaCl₂ and 0.1 mM MgCl₂ for 10 minutes at room temperature, followed by a permeabilization step with 0.2% Triton X-100 in the same buffer for 5 minutes . For paraffin embedded tissues use a pretreatment step of 15 minutes incubation in TRIS-EDTA buffer pH 9 in a microwave. Optimal antibody dilution should be determined by titration, recommended range is 1:50 - 1:100 for immunohistochemistry with avidin-biotinylated Horseradish peroxidase complex (ABC) as detection reagent, and 1:100 - 1:500 for immunoblotting applications.