

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Human CD2 FITC - CD20 PE, FITC/RPE NMB-0220S

Artikelname	Mouse anti Human CD2 FITC - CD20 PE, FITC/RPE
Artikelnummer	NMB-0220S
Hersteller Artikelnummer	0220S
Alternativnummer	NMB-0220S
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC
Spezies Reaktivität	Human
Immunogen	CD2=Derived from the hybridization of mouse Sp2/0 myeloma cells with spleen cells from BALB/c mice immunized with t lymphocytes activated by mixed lymphocyte culture.CD20=Derived from the hybridization of mouse Sp2/0 myeloma cells with spleen cells from BALB/c mice immunized with the LB lymphoblastoid cell line.
Konjugation	FITC/RPE
Produktbeschreibung	Identification of human T cells and subset of NK cells associated with the receptor for sheep erythrocytes rosettes expressing the 45-50,000 M.W. surface antigen. Identification of human T lymphocytes in multiple stages of T cell development, includin...
Konzentration	Titered for flow cytometry
Isotyp	IgG2a, IgG1

UniProt	P06729
Puffer	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Reinheit	Protein A/G Chromatography
Formulierung	Bi-Test (FITC/RPE) Reagent
Formel	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Anwendungsbeschreibung	PBMC: Add 10 µl of MAB/10⁶ PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8 C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature 20 C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.