

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Dog IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GAD/IGG(H+L)

Artikelname	Goat anti Dog IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAD/IGG(H+L)
Hersteller Artikelnummer	GAD/IgG(H+L)
Alternativnummer	NMB-GAD/IGG(H+L)
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Canine
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in dog serum, as t...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Quelle	Purified normal IgG isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 ml or equivalent against 120 ml antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 ml antiserum in a 3 mm diameter centre well and 2 ml serum samples (neat and diluted) in 2 mm diameter peripheral wells.