

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Horse IgG1(T), Clone: [Polyclonal], Monoclonal NMB-GAHO/IGG1(T)

Artikelname	Goat anti Horse IgG1(T), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GAHO/IGG1(T)
Hersteller Artikelnummer	GAHo/IgG1(T)
Alternativnummer	NMB-GAHO/IGG1(T)
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Equine
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG1 (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc subunit of subclass IgG1. It does not react with IgG/Fab fragments, IgM and IgA or any non-Ig protein in horse serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In preci...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protein
Quelle	Pools of purified normal IgG1 isolated from pooled serum from tetanus immunized horses. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protein
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.