

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IG(FC)/TRITC

Artikelname	Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-GARA/IG(FC)/TRITC
Hersteller Artikelnummer	GARa/Ig(Fc)/TRITC
Alternativnummer	NMB-GARA/IG(FC)/TRITC
Hersteller	NordicMubio
Wirt	Goat
Kategorie	Antikörper
Spezies Reaktivität	Rat
Konjugation	TRITC
Format	IgG
Spezifität	IgG+IgM+IgA (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc subunits of the major isotypes of the rat immunoglobulin system. No reaction is obtained with purified IgG/Fab or any non-Ig protein of rat serum, as tested by immunoelectrophoresis and double rad...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water.
Quelle	Purified polyclonal rat IgG, and homogenous IgA and IgM isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formulierung	TRITC
Formel	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.