

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Bovine IgG2 (subclass specific), Clone: [Polyclonal], Monoclonal NMB-RAB/IGG2

Artikelname	Rabbit anti Bovine IgG2 (subclass specific), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RAB/IGG2
Hersteller Artikelnummer	RAB/IgG2
Alternativnummer	NMB-RAB/IGG2
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Spezies Reaktivität	Bovine
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG2 (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc subunit of subclass IgG2. It does not react with other IgG1, IgG/Fab fragments, IgM and IgA or any non-Ig protein in bovine serum, as tested by immunoelectrophoresis and double radial immunodiffus...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign prote
Quelle	Pools of purified normal IgG2 isolated from pooled bovine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign prot
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.