

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Goat IgG1 IgG2 IgA IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAG/IG

Artikelname	Rabbit anti Goat IgG1 IgG2 IgA IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RAG/IG
Hersteller Artikelnummer	RAG/Ig
Alternativnummer	NMB-RAG/IG
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Spezies Reaktivität	Goat
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins. It does not react with any non-Ig protein in goat serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunoelectrophores...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rab
Quelle	Purified normal Ig fractions isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled ra
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.