

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit anti Human Factor VII, Clone: [Polyclonal], Monoclonal NMB-RAHU/FVII

|                          |                                                                                                                                                                                                                                                                 |
|--------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Rabbit anti Human Factor VII, Clone: [Polyclonal], Monoclonal                                                                                                                                                                                                   |
| Artikelnummer            | NMB-RAHU/FVII                                                                                                                                                                                                                                                   |
| Hersteller Artikelnummer | RAHu/FVII                                                                                                                                                                                                                                                       |
| Alternativnummer         | NMB-RAHU/FVII                                                                                                                                                                                                                                                   |
| Hersteller               | NordicMubio                                                                                                                                                                                                                                                     |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                          |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                      |
| Produktbeschreibung      | The defined antibody reactivity is restricted to Factor VII. To demonstrate the presence of FVII in normal plasma or serum by gel-immunodiffusion or immunoelectrophoresis a concentrate must be prepared, because the normal concentration is below the d...   |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                      |
| Klon-Bezeichnung         | [Polyclonal]                                                                                                                                                                                                                                                    |
| UniProt                  | <a href="#">P08709</a>                                                                                                                                                                                                                                          |
| Puffer                   | Delipidated, heat inactivated, lyophilized, stable whole serum. Sodium azide 1 mg/ml Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.                                    |
| Quelle                   | Plasma factor VII is ea vitamin K-dependent glycoprotein (MW 63,000) with an electrophoresis mobility in the beta region. It circulates in plasma in a semi-active form, even in the absence of tissue factor. Then procoagulant activity of FVII can be destro |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formel                 | <p>Delipidated, heat inactivated, lyophilized, stable whole serum.</p> <p>Sodium azide 1 mg/ml Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum.</p> <p>No foreign proteins added.</p>                                                                                                                                                                                                                                                                                                                                                                                                            |
| Anwendungsbeschreibung | <p>Precipitation assays. In immunoelectrophoresis in agarose-plates use 2 µl human plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl plasma samples (neat and serially diluted) in 2 mm diameter peripheral wells. In electroimmunodiffusion the antiserum concentration required in the gel is normally between 1 and 2%. The amount of Factor VII precipitated by 1 ml antiserum is approximately 100 U. One Unit of Factor VII is defined as the amount present in 1 ml normal plasma. On the average this corresponds to 1 µg/ml.</p> |