

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TACSTD2 / TROP2 (Epithelial Marker) Antibody, IgG1, Clone: [TACSTD2/2153], Mouse, Monoclonal NBT-4070-MSM3-P0

Artikelname	TACSTD2 / TROP2 (Epithelial Marker) Antibody, IgG1, Clone: [TACSTD2/2153], Mouse, Monoclonal
Artikelnummer	NBT-4070-MSM3-P0
Hersteller Artikelnummer	4070-MSM3-P0
Alternativnummer	NBT-4070-MSM3-P0-20,NBT-4070-MSM3-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Hamster, Human, Mouse, Rat
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Produktbeschreibung	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne...
Klonalität	Monoclonal
Klon-Bezeichnung	[TACSTD2/2153]
Molekulargewicht	40kDa

Isotyp	IgG1
NCBI	4070
UniProt	P09758
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a