

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HSP60 Antibody [LK1], Unconjugated, Mouse, Monoclonal PRS-33-141

| | |
|--------------------------|--|
| Artikelname | HSP60 Antibody [LK1], Unconjugated, Mouse, Monoclonal |
| Artikelnummer | PRS-33-141 |
| Hersteller Artikelnummer | 33-141 |
| Alternativnummer | PRS-33-141-100 |
| Hersteller | ProSci |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Bovine, Canine, Drosophila, Frog, Gallus, Hamster, Human, Monkey, Mouse, Porcine, Rabbit, Rat, Sheep |
| Immunogen | Recombinant human HSP60 was used as the immunogen for this antibody. |
| Konjugation | Unconjugated |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [LK1] |
| NCBI | 3329 |
| UniProt | P10809 |
| Puffer | PBS with 0.1 mg/ml BSA and 0.05% sodium azide |

| | |
|------------------------|--|
| Formulierung | Liquid |
| Application Verdünnung | Optimal dilutions for each application to be determined by the researcher |
| Anwendungsbeschreibung | <p>Flow Cytometry: 0.5-1 ug/million cells IF: 0.5-1 ug/ml WB: 0.25-0.5 ug/ml IHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1) Prediluted format : incubate for 30 min at RT (2) The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titered up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |