

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD63(LAMP3/968), CF405M conjugate, 0.1mg/mL, Clone: [LAMP3/968], Mouse, Monoclonal BOT-BNC050968-500**

|                          |   |
|--------------------------|---|
| Artikelname              | CD63(LAMP3/968), CF405M conjugate, 0.1mg/mL, Clone: [LAMP3/968], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC050968-500   |
| Hersteller Artikelnummer | BNC050968-500   |
| Alternativnummer         | BOT-BNC050968-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | Exosome Staining, FC, IF, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human full-length CD63 protein  |
| Konjugation              | CF405M  |
| Produktbeschreibung      | This MAb recognizes protein of 26 kDa-60 kDa, which is identified as CD63. Its epitope is different from that of MAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells an... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [LAMP3/968]   |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 26 kDa (core protein), 30-60 kDa (glycosylated)  |
| UniProt                | <a href="#">P08962</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |