

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CA19-9/Sialyl Lewisa (GI Tumor Marker) (CA19.9/1390R), CF405M conjugate, 0.1mg/mL, Clone: [CA19.9/1390R], Rabbit, Monoclonal BOT-BNC051390-100

| | |
|--------------------------|---|
| Artikelname | CA19-9/Sialyl Lewisa (GI Tumor Marker) (CA19.9/1390R), CF405M conjugate, 0.1mg/mL, Clone: [CA19.9/1390R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC051390-100 |
| Hersteller Artikelnummer | BNC051390-100 |
| Alternativnummer | BOT-BNC051390-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Purified human CA19-9 protein |
| Konjugation | CF405M |
| Produktbeschreibung | CA19-9, a carbohydrate epitope expressed on a high MW (>400 kDa) mucin glycoprotein, is a sialyl Lewisa structure which is synthesized from type 1 blood group precursor chains and is present in individuals expressing the Lewisa and/or Lewisb blood gr... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CA19.9/1390R] |

| | |
|------------------------|--|
| Molekulargewicht | >400 kDa |
| UniProt | Not Known |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence 0.5-1 ug/mL</p> <p>Immunohistology (formalin) 5-10 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |