

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytochrome C(CYCS/1010), CF640R conjugate, 0.1mg/mL, Clone: [CYCS/1010], Mouse, Monoclonal BOT-BNC401010-500**

Artikelname	Cytochrome C(CYCS/1010), CF640R conjugate, 0.1mg/mL, Clone: [CYCS/1010], Mouse, Monoclonal
Artikelnummer	BOT-BNC401010-500
Hersteller Artikelnummer	BNC401010-500
Alternativnummer	BOT-BNC401010-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC, WB
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant full-length human CYCS protein
Konjugation	CF640R
Produktbeschreibung	Cytochrome c is a well-characterized mobile electron transport protein that is essential to energy conversion in all aerobic organisms. In mammalian cells, this highly conserved protein is normally localized to the mitochondrial inter-membrane space....
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CYCS/1010]

Molekulargewicht	15 kDa
UniProt	<a href="#">P99999</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user