

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100B(S100B/1012), CF640R conjugate, 0.1mg/mL, Clone: [S100B/1012], Mouse, Monoclonal BOT-BNC401012-500

| | |
|--------------------------|---|
| Artikelname | S100B(S100B/1012), CF640R conjugate, 0.1mg/mL, Clone: [S100B/1012], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401012-500 |
| Hersteller Artikelnummer | BNC401012-500 |
| Alternativnummer | BOT-BNC401012-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC |
| Spezies Reaktivität | Bovine, Human, Mouse, Rat |
| Immunogen | Recombinant full-length human S100B protein |
| Konjugation | CF640R |
| Produktbeschreibung | S100 belongs to the family of calcium binding proteins. S100A and S100B proteins are two members of the S100 family. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epi... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [S100B/1012] |

| | |
|------------------------|---|
| Molekulargewicht | 10-12 kDa |
| UniProt | P04271 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |