

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD50 / ICAM3(ICO-60), CF640R conjugate, 0.1mg/mL, Clone: [ICO-60], Mouse, Monoclonal BOT-BNC401040-500**

|                          |   |
|--------------------------|---|
| Artikelname              | CD50 / ICAM3(ICO-60), CF640R conjugate, 0.1mg/mL, Clone: [ICO-60], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC401040-500   |
| Hersteller Artikelnummer | BNC401040-500   |
| Alternativnummer         | BOT-BNC401040-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, Functional Studies  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human ICAM3   |
| Konjugation              | CF640R  |
| Produktbeschreibung      | Recognizes the N-glycosylated glycoprotein of 120 kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3. CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhesion. It is expressed on thymocytes and ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [ICO-60]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 110-160 kDa   |
| UniProt                | <a href="#">P32942</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |