

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Chromogranin A(CGA414 + CGA/777 + CGA/798), CF640R conjugate, 0.1mg/mL, Clone: [CGA414 CGA/777 CGA/798], Mouse, Monoclonal BOT-BNC401060-100

Artikelname	Chromogranin A(CGA414 + CGA/777 + CGA/798), CF640R conjugate, 0.1mg/mL, Clone: [CGA414 CGA/777 CGA/798], Mouse, Monoclonal
Artikelnummer	BOT-BNC401060-100
Hersteller Artikelnummer	BNC401060-100
Alternativnummer	BOT-BNC401060-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Monkey, Mouse, Porcine, Rat
Immunogen	Recombinant human chromogranin A protein
Konjugation	CF640R
Produktbeschreibung	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CGA414 CGA/777 CGA/798]

Molekulargewicht	68-75 kDa
UniProt	P10645
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user