

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **S100A9(S100A9/1075), CF640R conjugate, 0.1mg/mL, Clone: [S100A9/1075], Mouse, Monoclonal BOT-BNC401075-500**

|                          |   |
|--------------------------|---|
| Artikelname              | S100A9(S100A9/1075), CF640R conjugate, 0.1mg/mL, Clone: [S100A9/1075], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC401075-500   |
| Hersteller Artikelnummer | BNC401075-500   |
| Alternativnummer         | BOT-BNC401075-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human, Rat  |
| Immunogen                | Recombinant human S100A9 protein  |
| Konjugation              | CF640R  |
| Produktbeschreibung      | This MAb stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief ce... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [S100A9/1075]   |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 14 kDa   |
| UniProt                | <a href="#">P06702</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT Flow Cytometry 0.5-1 ug/million cells/0.1 mL |