

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD51 / Integrin V(ITGAV/1610), CF640R conjugate, 0.1mg/mL, Clone: [ITGAV/1610], Mouse, Monoclonal BOT-BNC401610-100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | CD51 / Integrin V(ITGAV/1610), CF640R conjugate, 0.1mg/mL, Clone: [ITGAV/1610], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401610-100 |
| Hersteller Artikelnummer | BNC401610-100 |
| Alternativnummer | BOT-BNC401610-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length integrin alpha V beta 1 protein |
| Konjugation | CF640R |
| Produktbeschreibung | Integrins are heterodimers composed of noncovalently associated transmembrane both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ITGAV/1610] |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht | 150-160 kDa under non-reducing conditions. Under reducing conditions, it is cleaved into 2 bands of about 125-130 kDa 20-25 kDa. |
| UniProt | P06756 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |