

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1644), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [CFTR/1644], Mouse, Monoclonal BOT-BNC401644-100

| | |
|--------------------------|---|
| Artikelname | CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1644), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [CFTR/1644], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401644-100 |
| Hersteller Artikelnummer | BNC401644-100 |
| Alternativnummer | BOT-BNC401644-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human CFTR fragment (aa258-385) (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structu... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [CFTR/1644] |
| Molekulargewicht | 165-170 kDa |
| Isotyp | IgG1 |
| UniProt | P13569 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |