

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD79a (HM57), CF640R conjugate, 0.1mg/mL, Clone: [HM57], Mouse, Monoclonal BOT-BNC401678-500**

Artikelname	CD79a (HM57), CF640R conjugate, 0.1mg/mL, Clone: [HM57], Mouse, Monoclonal
Artikelnummer	BOT-BNC401678-500
Hersteller Artikelnummer	BNC401678-500
Alternativnummer	BOT-BNC401678-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Bovine, Human, Monkey, Mouse, Porcine, Rat
Immunogen	A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein.
Konjugation	CF640R
Produktbeschreibung	A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[HM57]
Molekulargewicht	44 kDa
UniProt	<a href="#">P11912</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user