

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p63 (Squamous, Basal & Myoepithelial Cell Marker), CF640R conjugate, 0.1mg/mL, Rabbit BOT-BNC401684-100

| | |
|--------------------------|---|
| Artikelname | p63 (Squamous, Basal & Myoepithelial Cell Marker), CF640R conjugate, 0.1mg/mL, Rabbit |
| Artikelnummer | BOT-BNC401684-100 |
| Hersteller Artikelnummer | BNC401684-100 |
| Alternativnummer | BOT-BNC401684-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human p63 protein fragment (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal... |
| Konzentration | 0.1 mg/mL |
| Molekulargewicht | 63 kDa |

| | |
|------------------------|--|
| UniProt | Q9H3D4 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |