

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD44v4 (Marker of Tumor Metastasis) (CD44v4/1700R), CF640R conjugate, 0.1mg/mL, Clone: [CD44v4/1700R], Rabbit, Monoclonal BOT-BNC401700-500

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | CD44v4 (Marker of Tumor Metastasis) (CD44v4/1700R), CF640R conjugate, 0.1mg/mL, Clone: [CD44v4/1700R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC401700-500 |
| Hersteller Artikelnummer | BNC401700-500 |
| Alternativnummer | BOT-BNC401700-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment corresponding to the v4 domain of human CD44 (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes an epitope encoded by exon v4 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemop... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Klon-Bezeichnung | [CD44v4/1700R] |
| Molekulargewicht | 80-95 kDa |
| UniProt | P16070 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |