

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Chromogranin A / CHGA (Neuroendocrine Marker), CF640R conjugate, 0.1mg/mL, Rabbit BOT-BNC401732-500**

Artikelname	Chromogranin A / CHGA (Neuroendocrine Marker), CF640R conjugate, 0.1mg/mL, Rabbit
Artikelnummer	BOT-BNC401732-500
Hersteller Artikelnummer	BNC401732-500
Alternativnummer	BOT-BNC401732-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length Chromogranin A protein
Konjugation	CF640R
Produktbeschreibung	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para...
Konzentration	0.1 mg/mL
Molekulargewicht	68-75 kDa
UniProt	<a href="#">P10645</a>

Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user