

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD34 (Hematopoietic Stem Cell & Endothelial Marker) (HPCA1/1806R), CF640R conjugate, 0.1mg/mL, Clone: [HPCA1/1806R], Rabbit, Monoclonal BOT-BNC401806-100

| | |
|--------------------------|---|
| Artikelname | CD34 (Hematopoietic Stem Cell & Endothelial Marker) (HPCA1/1806R), CF640R conjugate, 0.1mg/mL, Clone: [HPCA1/1806R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC401806-100 |
| Hersteller Artikelnummer | BNC401806-100 |
| Alternativnummer | BOT-BNC401806-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant full-length human HPCA1 protein |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes a transmembrane, heavily glycosylated protein of 90-120 kDa, which is identified as CD34. Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as li... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [HPCA1/1806R] |
| Molekulargewicht | 90-110 kDa |
| UniProt | P28906 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Immunohistology (formalin) 2-4 ug/mL Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |