

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Beta-2 Microglobulin (Renal Failure & Tumor Marker) (B2M/1857R), CF640R conjugate, 0.1mg/mL, Clone: [B2M/1857R], Rabbit, Monoclonal BOT-BNC401857-500

Artikelname	Beta-2 Microglobulin (Renal Failure & Tumor Marker) (B2M/1857R), CF640R conjugate, 0.1mg/mL, Clone: [B2M/1857R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC401857-500
Hersteller Artikelnummer	BNC401857-500
Alternativnummer	BOT-BNC401857-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Primate
Immunogen	Recombinant full-length human B2M protein
Konjugation	CF640R
Produktbeschreibung	Beta-2 microglobulin is a component of the class I major histocompatibility complex (MHC I) where it functions in antigen processing and presentation. Involved in the presentation of peptide antigens to the immune system. Human Beta-2 microglobulin i...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[B2M/1857R]
Molekulargewicht	12 kDa
UniProt	P61769
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user