

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Kappa Light Chain / IGKC (B-Cell Marker) (KLC2289R), CF647 conjugate, 0.1mg/mL, Clone: [KLC2289R], Rabbit, Monoclonal BOT-BNC472289-100

Artikelname	Human Kappa Light Chain / IGKC (B-Cell Marker) (KLC2289R), CF647 conjugate, 0.1mg/mL, Clone: [KLC2289R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472289-100
Hersteller Artikelnummer	BNC472289-100
Alternativnummer	BOT-BNC472289-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Ig kappa chain (IGKC) protein
Konjugation	CF647
Produktbeschreibung	This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. It recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects al...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[KLC2289R]

Molekulargewicht	~22.5 kDa
UniProt	P01601
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user