

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Thrombomodulin / CD141 (Endothelial Cell Marker) (rTHBD/1591), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [rTHBD/1591], Mouse, Monoclonal BOT-BNC472332-100

| | |
|--------------------------|---|
| Artikelname | Thrombomodulin / CD141 (Endothelial Cell Marker) (rTHBD/1591), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [rTHBD/1591], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC472332-100 |
| Hersteller Artikelnummer | BNC472332-100 |
| Alternativnummer | BOT-BNC472332-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 69-194) of human Thrombomodulin (CD141) protein (exact sequence is proprietary) |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes a protein of 75 kDa, identified as Thrombomodulin. Thrombomodulin is a transmembrane glycoprotein with natural anticoagulant properties. It is normally expressed by a restricted number of cells, such as endothelial and mesoth... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [rTHBD/1591] |
| Molekulargewicht | 100 kDa |
| Isotyp | IgG1 |
| UniProt | P07204 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |