

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TRAcP (Tartrate-Resistant Acid Phosphatase) (Hairy Cell Leukemia Marker) (ACP5/2336R), CF647 conjugate, 0.1mg/mL, Clone: [ACP5/2336R], Rabbit, Monoclonal BOT-BNC472336-100

| | |
|--------------------------|---|
| Artikelname | TRAcP (Tartrate-Resistant Acid Phosphatase) (Hairy Cell Leukemia Marker) (ACP5/2336R), CF647 conjugate, 0.1mg/mL, Clone: [ACP5/2336R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC472336-100 |
| Hersteller Artikelnummer | BNC472336-100 |
| Alternativnummer | BOT-BNC472336-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Recombinant full-length human ACP5 protein |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes a protein of 35 kDa, which is identified as tartrate-resistant acid phosphatase (TRAcP). It exists as two isoforms (5a and 5b). This MAb reacts with both the isoforms. Serum TRAcP 5a is secreted by macrophages and dendritic c... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [ACP5/2336R] |
| Molekulargewicht | 35 kDa |
| UniProt | P13686 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |