

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD44v9 (Marker of Tumor Metastasis) (CD44v9/2344R), CF647 conjugate, 0.1mg/mL, Clone: [CD44v9/2344R], Rabbit, Monoclonal BOT-BNC472344-100

| | |
|--------------------------|---|
| Artikelname | CD44v9 (Marker of Tumor Metastasis) (CD44v9/2344R), CF647 conjugate, 0.1mg/mL, Clone: [CD44v9/2344R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC472344-100 |
| Hersteller Artikelnummer | BNC472344-100 |
| Alternativnummer | BOT-BNC472344-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment corresponding to the v9 domain of human CD44 |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes an epitope encoded by exon v4 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemop... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [CD44v9/2344R] |
| Molekulargewicht | 80-95 kDa |
| UniProt | P16070 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |