

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Epithelial Marker) (rEGP40/1110), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [rEGP40/1110], Mouse, Monoclonal BOT-BNC472347-500

Artikelname	Ep-CAM / CD326 (Epithelial Marker) (rEGP40/1110), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [rEGP40/1110], Mouse, Monoclonal
Artikelnummer	BOT-BNC472347-500
Hersteller Artikelnummer	BNC472347-500
Alternativnummer	BOT-BNC472347-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant human EpCAM fragment from the cytoplasmic domain (around aa 280-350) (exact sequence is proprietary)
Konjugation	CF647
Produktbeschreibung	EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majo...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rEGP40/1110]
Molekulargewicht	40-43 kDa
Isotyp	IgG1
UniProt	P16422
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user