

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC16 / CA125 (Ovarian Carcinoma Marker) (OCA125/2349R), CF647 conjugate, 0.1mg/mL, Clone: [OCA125/2349R], Rabbit, Monoclonal BOT-BNC472349-500

Artikelname	MUC16 / CA125 (Ovarian Carcinoma Marker) (OCA125/2349R), CF647 conjugate, 0.1mg/mL, Clone: [OCA125/2349R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472349-500
Hersteller Artikelnummer	BNC472349-500
Alternativnummer	BOT-BNC472349-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human MUC16 protein
Konjugation	CF647
Produktbeschreibung	The mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats (VNTRs). Membrane-associated and secretory Mucins are high molecular weight glycoproteins of the glycocalyx (po...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[OCA125/2349R]
Molekulargewicht	>2,000 kDa
UniProt	Q8WXI7
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user