

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF647 conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal BOT-BNC472368-100**

Artikelname	Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF647 conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal
Artikelnummer	BOT-BNC472368-100
Hersteller Artikelnummer	BNC472368-100
Alternativnummer	BOT-BNC472368-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Cytokeratin 6A (KRT6A) protein
Konjugation	CF647
Produktbeschreibung	This MAb recognizes a protein of 56 kDa, identified as cytokeratin 6A (KRT6A). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns. Cytokeratin 6A is the dominant...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[KRT6A/2368]

Molekulargewicht	56 kDa
UniProt	<a href="#">P02538</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user