

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **BOB.1 (B-Cell Marker)(BOB1/2422), CF647 conjugate, 0.1mg/mL, Clone: [BOB1/2422], Mouse, Monoclonal BOT-BNC472422-500**

Artikelname	BOB.1 (B-Cell Marker)(BOB1/2422), CF647 conjugate, 0.1mg/mL, Clone: [BOB1/2422], Mouse, Monoclonal
Artikelnummer	BOT-BNC472422-500
Hersteller Artikelnummer	BNC472422-500
Alternativnummer	BOT-BNC472422-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa 148-255) of human BOB1 (POU2AF1) protein (exact sequence is proprietary)
Konjugation	CF647
Produktbeschreibung	BOB-1 is a transcriptional coactivator that specifically associates with either POU2F1/OCT1 or POU2F2/OCT2, boosting the promoter activity of those complexes. It is essential for the response of B-cells to antigens, required for the formation of germ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[BOB1/2422]
Molekulargewicht	35 kDa
UniProt	<a href="#">Q16633</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95AC followed by cooling at RT for 20 minutes Immunohistology (formalin): 0.1-0.2 ug/mL for 30 minutes at RT Western Blot 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user