

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

BOB.1 (B-Cell Marker) (BOB1/2425), CF647 conjugate, 0.1mg/mL, Clone: [BOB1/2425], Mouse, Monoclonal BOT-BNC472425-100

Artikelname	BOB.1 (B-Cell Marker) (BOB1/2425), CF647 conjugate, 0.1mg/mL, Clone: [BOB1/2425], Mouse, Monoclonal
Artikelnummer	BOT-BNC472425-100
Hersteller Artikelnummer	BNC472425-100
Alternativnummer	BOT-BNC472425-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human BOB1 (POU2AF1) protein (around aa 148-255) (exact sequence is proprietary)
Konjugation	CF647
Produktbeschreibung	BOB-1 is a transcriptional coactivator that specifically associates with either POU2F1/OCT1 or POU2F2/OCT2, boosting the promoter activity of those complexes. It is essential for the response of B-cells to antigens, required for the formation of germ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[BOB1/2425]
Molekulargewicht	35 kDa
UniProt	Q16633
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user