

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2428), CF647 conjugate, 0.1mg/mL, Clone: [TP63/2428], Mouse, Monoclonal BOT-BNC472428-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2428), CF647 conjugate, 0.1mg/mL, Clone: [TP63/2428], Mouse, Monoclonal                                                                                                                               |
| Artikelnummer            | BOT-BNC472428-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC472428-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC472428-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IHC                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human p63 protein                                                                                                                                                                                                                     |
| Konjugation              | CF647                                                                                                                                                                                                                                                         |
| Produktbeschreibung      | p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [TP63/2428]                                                                                                                                                                                                                                                   |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 63 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| UniProt                | <a href="#">Q9H3D4</a>                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |