

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD3e (T-Cell Marker) (C3e/2479), CF647 conjugate, 0.1mg/mL, Clone: [C3e/2479], Mouse, Monoclonal BOT-BNC472479-100

Artikelname	CD3e (T-Cell Marker) (C3e/2479), CF647 conjugate, 0.1mg/mL, Clone: [C3e/2479], Mouse, Monoclonal
Artikelnummer	BOT-BNC472479-100
Hersteller Artikelnummer	BNC472479-100
Alternativnummer	BOT-BNC472479-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	WB
Spezies Reaktivität	Human
Immunogen	Recombinant human CD3e extracellular domain fragment (around aa 23-119) (exact sequence is proprietary)
Konjugation	CF647
Produktbeschreibung	The immunogen for this Mab is derived from the extracellular domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closel...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[C3e/2479]
Molekulargewicht	20 kDa
UniProt	P07766
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user