

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MUC18 / CD146 / MCAM (Melanoma Cell Adhesion Molecule) (rMUC18/1130), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rMUC18/1130], Mouse, Monoclonal BOT-BNC682302-500**

Artikelname	MUC18 / CD146 / MCAM (Melanoma Cell Adhesion Molecule) (rMUC18/1130), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rMUC18/1130], Mouse, Monoclonal
Artikelnummer	BOT-BNC682302-500
Hersteller Artikelnummer	BNC682302-500
Alternativnummer	BOT-BNC682302-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human MUC18 protein
Konjugation	CF568
Produktbeschreibung	Mel-CAM belongs to the immunoglobulin superfamily and functions as a Ca <sup>2+</sup> -independent cell adhesion molecule. Mel-CAM expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not norm...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rMUC18/1130]
Molekulargewicht	130 kDa
Isotyp	IgG1
UniProt	<a href="#">P43121</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA pH 8.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user