

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TRAcP (Tartrate-Resistant Acid Phosphatase) (Hairy Cell Leukemia Marker) (rACP5/1070), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rACP5/1070], Mouse, Monoclonal BOT-BNC682337-100

Artikelname	TRAcP (Tartrate-Resistant Acid Phosphatase) (Hairy Cell Leukemia Marker) (rACP5/1070), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rACP5/1070], Mouse, Monoclonal
Artikelnummer	BOT-BNC682337-100
Hersteller Artikelnummer	BNC682337-100
Alternativnummer	BOT-BNC682337-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant full-length human ACP5 protein
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 35 kDa, which is identified as tartrate-resistant acid phosphatase (TRAcP). It exists as two isoforms (5a and 5b). This MAb reacts with both the isoforms. Serum TRAcP 5a is secreted by macrophages and dendritic c...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rACP5/1070]
Molekulargewicht	35 kDa
Isotyp	IgG1
UniProt	P13686
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user