

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/2338R), CF568 conjugate, 0.1mg/mL, Clone: [FTL/2338R], Rabbit, Monoclonal BOT-BNC682338-500**

Artikelname	Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/2338R), CF568 conjugate, 0.1mg/mL, Clone: [FTL/2338R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC682338-500
Hersteller Artikelnummer	BNC682338-500
Alternativnummer	BOT-BNC682338-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human FTL protein fragment (around aa 38-165) (exact sequence is proprietary)
Konjugation	CF568
Produktbeschreibung	Mammalian ferritins consist of 24 subunits made up of 2 types of polypeptide chains, ferritin heavy chain and ferritin light chain. Ferritin heavy chains catalyze the first step in iron storage, the oxidation of Fe (II), whereas ferritin light chains...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[FTL/2338R]
Molekulargewicht	19-25 kDa
UniProt	<a href="#">P02792</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Western Blotting 0.5-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user