

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker) (TYRP1/2340R), CF568 conjugate, 0.1mg/mL, Clone: [TYRP1/2340R], Rabbit, Monoclonal BOT-BNC682340-100

| | |
|--------------------------|---|
| Artikelname | Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker) (TYRP1/2340R), CF568 conjugate, 0.1mg/mL, Clone: [TYRP1/2340R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC682340-100 |
| Hersteller Artikelnummer | BNC682340-100 |
| Alternativnummer | BOT-BNC682340-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human TYRP1 protein (around aa 257-377) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | This antibody reacts with a 75 kDa melanocyte-specific gene product, identified as Tyrosinase-Related Protein-1 (TRP-1). It is involved in melanin synthesis. TRP1 is present on the melanosomal membranes of melanoma, normal melanocytes and nevi. Recen... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [TYRP1/2340R] |
| Molekulargewicht | 75 kDa |
| UniProt | P17643 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |