

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD269 / TNFRSF17 / BCMA (B-Cell Maturation Protein) (BCMA/2366), CF568 conjugate, 0.1mg/mL, Clone: [BCMA/2366], Mouse, Monoclonal BOT-BNC682366-100**

Artikelname	CD269 / TNFRSF17 / BCMA (B-Cell Maturation Protein) (BCMA/2366), CF568 conjugate, 0.1mg/mL, Clone: [BCMA/2366], Mouse, Monoclonal
Artikelnummer	BOT-BNC682366-100
Hersteller Artikelnummer	BNC682366-100
Alternativnummer	BOT-BNC682366-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human CD269 protein fragment (around aa 78-184) (exact sequence is proprietary)
Konjugation	CF568
Produktbeschreibung	The B cell maturation protein (BCMA) is a type I integral membrane protein that belongs to the tumor necrosis factor receptor (TNF-R) superfamily. It is expressed as a 184 amino acid peptide that is expressed only in mature B-lymphocytes and is locat...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[BCMA/2366]
Molekulargewicht	20 kDa
UniProt	<a href="#">Q02223</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: For coating order antibody without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user