

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF568 conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal BOT-BNC682368-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF568 conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC682368-100   |
| Hersteller Artikelnummer | BNC682368-100   |
| Alternativnummer         | BOT-BNC682368-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human Cytokeratin 6A (KRT6A) protein  |
| Konjugation              | CF568   |
| Produktbeschreibung      | This MAb recognizes a protein of 56 kDa, identified as cytokeratin 6A (KRT6A). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns. Cytokeratin 6A is the dominant... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [KRT6A/2368]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 56 kDa  |
| UniProt                | <a href="#">P02538</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |