

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Histone H1 (Nuclear Marker) (r1415-1), 0.2mg/mL, Clone: [r1415-1], Mouse, Monoclonal BOT-BNUB1797-100

Artikelname	Histone H1 (Nuclear Marker) (r1415-1), 0.2mg/mL, Clone: [r1415-1], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1797-100
Hersteller Artikelnummer	BNUB1797-100
Alternativnummer	BOT-BNUB1797-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Nuclei of human leukemia biopsy cells
Produktbeschreibung	Eukaryotic histones are basic and water-soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed supealpha-helicalturn sequentially to form chromosomal fiber. Two molecules of each of...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[r1415-1]
Molekulargewicht	~30 kDa

UniProt	Multiple
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user