

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) (rKi-1/779), 0.2mg/mL, Clone: [rKi-1/779], Mouse, Monoclonal BOT-BNUB1828-500

Artikelname	CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) (rKi-1/779), 0.2mg/mL, Clone: [rKi-1/779], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1828-500
Hersteller Artikelnummer	BNUB1828-500
Alternativnummer	BOT-BNUB1828-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human TNFRSF8 protein
Produktbeschreibung	This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[rKi-1/779]
Molekulargewicht	105-120 kDa

UniProt	P28908
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA pH 7.5-8.5 Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user