

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 7 (Glandular and Transitional Epithelial Marker) (rOV-TL12/30), 0.2mg/mL, IgG1, Clone: [rOV-TL12/30], Mouse, Monoclonal BOT-BNUB1841-500

Artikelname	Cytokeratin 7 (Glandular and Transitional Epithelial Marker) (rOV-TL12/30), 0.2mg/mL, IgG1, Clone: [rOV-TL12/30], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1841-500
Hersteller Artikelnummer	BNUB1841-500
Alternativnummer	BOT-BNUB1841-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC, WB
Spezies Reaktivität	Human
Immunogen	OTN 11 ovarian carcinoma cell line
Produktbeschreibung	This antibody recognizes an intermediate filament protein (IFP) of 55 kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[rOV-TL12/30]
Molekulargewicht	55 kDa

Isotyp	IgG1
UniProt	P08729
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user