

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Wilm s Tumor 1 (WT1) (Wilm s Tumor & Mesothelial Marker) (rWT1/857), 0.2mg/mL, Clone: [rWT1/857], Mouse, Monoclonal BOT-BNUB1856-500

Artikelname	Wilm s Tumor 1 (WT1) (Wilm s Tumor & Mesothelial Marker) (rWT1/857), 0.2mg/mL, Clone: [rWT1/857], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1856-500
Hersteller Artikelnummer	BNUB1856-500
Alternativnummer	BOT-BNUB1856-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human WT1 protein
Produktbeschreibung	This antibody recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subse...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[rWT1/857]
Molekulargewicht	47-55 kDa

UniProt	P19544
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user