

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD8A (Cytotoxic-&Suppressor T-Cell Marker) (rC8/468), 0.2mg/mL, Clone: [rC8/468], Mouse, Monoclonal BOT-BNUB1867-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD8A (Cytotoxic-&Suppressor T-Cell Marker) (rC8/468), 0.2mg/mL, Clone: [rC8/468], Mouse, Monoclonal                                                                                                                                                           |
| Artikelnummer            | BOT-BNUB1867-100                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB1867-100                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB1867-100-100UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human CD8a protein                                                                                                                                                                                                                    |
| Produktbeschreibung      | CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC ... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [rC8/468]                                                                                                                                                                                                                                                     |
| Molekulargewicht         | 32 kDa                                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                                                                                                                    |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt                | <a href="#">P01732</a>                                                                                                                                                                                                                                                                                                                             |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                        |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                             |
| Anwendungsbeschreibung | Not recommended for flow cytometry Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |