

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD8A (Cytotoxic-&Suppressor T-Cell Marker) (rC8/468), 0.2mg/mL, Clone: [rC8/468], Mouse, Monoclonal BOT-BNUB1867-500

Artikelname	CD8A (Cytotoxic-&Suppressor T-Cell Marker) (rC8/468), 0.2mg/mL, Clone: [rC8/468], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1867-500
Hersteller Artikelnummer	BNUB1867-500
Alternativnummer	BOT-BNUB1867-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CD8a protein
Produktbeschreibung	CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[rC8/468]
Molekulargewicht	32 kDa

UniProt	P01732
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Not recommended for flow cytometry Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user