

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TLE1 (Synovial Sarcoma Marker) (TLE1/2051), 0.2mg/mL, Clone: [TLE1/2051], Mouse, Monoclonal BOT-BNUB2051-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | TLE1 (Synovial Sarcoma Marker) (TLE1/2051), 0.2mg/mL, Clone: [TLE1/2051], Mouse, Monoclonal                                                                                                                                                                   |
| Artikelnummer            | BOT-BNUB2051-100                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB2051-100                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB2051-100-100UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC, WB                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human TLE1 fragment (aa 175-338) (exact sequence is proprietary)                                                                                                                                                                                  |
| Produktbeschreibung      | Key players in the Notch pathway are the TLE genes, which are human homologs of the Drosophila groucho gene. Groucho is a transcriptional repressor that plays a key role in neurogenesis, segmentation and sex determination. Transducin-like enhancer pr... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [TLE1/2051]                                                                                                                                                                                                                                                   |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 83 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| UniProt                | <a href="#">Q04724</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |