

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Carcinoembryonic Antigen (CEA) / CD66 (rC66/1009), 0.2mg/mL, IgG1, Clone: [rC66/1009], Mouse, Monoclonal BOT-BNUB2054-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Carcinoembryonic Antigen (CEA) / CD66 (rC66/1009), 0.2mg/mL, IgG1, Clone: [rC66/1009], Mouse, Monoclonal                                                                                                                                                      |
| Artikelnummer            | BOT-BNUB2054-100                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB2054-100                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB2054-100-100UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human CEA protein                                                                                                                                                                                                                     |
| Produktbeschreibung      | This antibody recognizes proteins of 80-200 kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MA... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [rC66/1009]                                                                                                                                                                                                                                                   |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 80-200 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Isotyp                 | IgG1                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| UniProt                | <a href="#">P06731</a>                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |